ANOGEN - A Division of YES Biotech Laboratories Ltd. 2355 Derry Road East, Unit 23, Mississauga, ON, Canada L55 1V6 • Tel: (905) 677-9221 • Fax: (905) 677-0023

Mouse Anti-Human IL-16 Monoclonal Antibody Datasheet

D-C40089H Mouse monoclonal antibody to human interleukin-16 (IL-16) Protein G affinity purified Primary antibody Human IL-16 Purified recombinant human IL-16 Sp2/0-Ag14 Reactive with natural and recombinant	Quantity: 0.5 mg Reconstitution: Storage: Research Area: Background:	Double distilled water is recommended to adjust the final concentration to 1.00mg/mL. Store at -20°C Cytokine Interleukin 16 (IL-16) is a pleiotropic cytokine, possessing immunomodulation function to allergic inflammatory diseases such as Crohn's disease and
interleukin-16 (IL-16) Protein G affinity purified Primary antibody Human IL-16 Purified recombinant human IL-16 Sp2/0-Ag14	Storage: Research Area:	to adjust the final concentration to 1.00mg/mL. Store at -20°C Cytokine Interleukin 16 (IL-16) is a pleiotropic cytokine, possessing immunomodulation function to allergic inflammatory diseases such as Crohn's disease and
Human IL-16 Purified recombinant human IL-16 Sp2/0-Ag14	Research Area:	Cytokine Interleukin 16 (IL-16) is a pleiotropic cytokine, possessing immunomodulation function to allergic inflammatory diseases such as Crohn's disease and
Purified recombinant human IL-16 Sp2/0-Ag14	Background:	cytokine, possessing immunomodulation function to allergic inflammatory diseases such as Crohn's disease and
human IL-16 Human, others not tested		ulcerative colitis. IL-16 utilizes CD4 as its primary receptor. IL-16 is a chemo- attractant to CD4+ T helper 1 cells. To CD4+ T helper 2 cells, IL-16 has an inhibitory effect on the antigen- mediated CD3/T cell receptor activation.
mAb 210 exhibits no detectable cross- reaction with human serum albumin, bovine serum albumin and other human cytokines or growth factors tested, such as IL-1 β , IL-8, IL-7 MCAF, TGF- β 1, EGF, bFGF, GM-CSF, M-CSF, TNF- α . Mouse, IgG2a Kappa Lyophilized from a solution in 0.01M	Applications: References:	 IL-16 has been shown to suppress HIV infection in lymphocytes and monocytes. ELISA If research is published using this product, please inform Anogen in order to cite the reference on this datasheet. Anogen will provide one unit of product
rreb cy a: b N Ly	Ab 210 exhibits no detectable cross- eaction with human serum albumin, ovine serum albumin and other human ytokines or growth factors tested, such s IL-1β, IL-8, IL-7 MCAF, TGF-β1, EGF, FGF, GM-CSF, M-CSF, TNF- α .	Ab 210 exhibits no detectable cross- eaction with human serum albumin, ovine serum albumin and other human ytokines or growth factors tested, such s IL-1β, IL-8, IL-7 MCAF, TGF-β1, EGF, FGF, GM-CSF, M-CSF, TNF- α .Applications: References:Nouse, IgG2a Kappa yophilized from a solution in 0.01M α α

This product is for LABORATORY RESEARCH USE and further manufacture ONLY, and cannot be administrated to human and animals for use in diagnostic and therapeutic procedures. Manufactured by ANOGEN - A Division of YES Biotech Laboratories Ltd.

Page 1 of 1 page(s)

To place order, please contact us by phone, fax or by email: <u>info@anogen.ca</u>, or with our secure online store: <u>www.anogen.net</u> S7.5 (02)